

Scoring Key

Symptom	Score					
	-1	0	1	2	3	4
Epistaxis	--	No or trivial (less than 5)	> 5 or more than 10'	Consultation only	Packing or cauterization or antifibrinolytic	Blood transfusion or replacement therapy or desmopressin
Cutaneous	--	No or trivial (< 1cm)	> 1 cm and no trauma	Consultation only	--	--
Bleeding from minor wounds	--	No or trivial (less than 5)	> 5 or more than 5'	Consultation only	Surgical hemostasis	Blood transfusion or replacement therapy or desmopressin
Oral cavity	--	No	Referred at least one	Consultation only	Surgical hemostasis or antifibrinolytic	Blood transfusion or replacement therapy or desmopressin
Gastrointestinal bleeding	--	No	Associated with ulcer, portal hypertension, hemorrhoids, angiodysplasia	Spontaneous	Surgical hemostasis, blood transfusion, replacement therapy, desmopressin, antifibrinolytic	--
Tooth extraction	No bleeding in at least 2 extractions	None done or no bleeding in 1 extraction	Reported, no consultation	Consultation only	Resuturing or packing	Blood transfusion or replacement therapy or desmopressin
Surgery	No bleeding in at least 2 surgeries	None done or no bleeding in 1 surgery	Reported, no consultation	Consultation only	Surgical hemostasis or antifibrinolytic	Blood transfusion or replacement therapy or desmopressin
Menorrhagia	--	No	Consultation only	Antifibrinolytics, pill use	Dilation & curettage, iron therapy, ablation	Blood transfusion or replacement therapy or desmopressin or hysterectomy
Postpartum hemorrhage	No bleeding in at least 2 deliveries	None done or no bleeding in 1 surgery	Consultation only	Dilation & curettage, iron therapy, antifibrinolytics	Blood transfusion or replacement therapy or desmopressin	Hysterectomy
Muscle hematomas	--	Never	Post trauma, no therapy	Spontaneous, no therapy	Spontaneous or traumatic, requiring desmopressin or replacement therapy	Spontaneous or traumatic, requiring surgical intervention or blood transfusion
Hemarthrosis	--	Never	Post trauma, no therapy	Spontaneous, no therapy	Spontaneous or traumatic, requiring desmopressin or replacement therapy	Spontaneous or traumatic, requiring surgical intervention or blood transfusion
Central nervous system bleeding	--	Never	--	--	Subdural, any intervention	Intracerebral, any intervention

Scoring Key: A few small changes in wording from the original MCMDM-1 scoring key are indicated in the boxes. For tooth extraction and surgery a score of 1 was previously for "Referred in <25% of cases" and a score of 2 was previously for "Referred in >25% of cases". For the menorrhagia category we added ablation therapy (which was not included in the original) at the same level as D&C and iron therapy.