

# Factor FVIII Deficiency - 2020

## Contents

<b>Factor FVIII Deficiency - Statistics Canada age groups</b>	<b>2</b>
<b>Factor FVIII Deficiency - Patients with an active inhibitor, Statistics Canada age groups</b>	<b>3</b>
<b>Factor FVIII Deficiency - Pediatric and Adult age groups</b>	<b>4</b>

Note: Prior national demographic reports were produced from the Canadian Hemophilia Registry. Beginning with 2018, these reports are now produced from the Canadian Blood Disorders Registry, which covers all of Canada except British Columbia. Numbers for British Columbia were retrieved separately and added to the CBDR totals.

## Factor FVIII Deficiency - Statistics Canada age groups

Coverage: All of Canada

Data source: CBDR, BC iChip

Year: 2020

Note: Inhibitor patients are included in this table

Severity	GenderCode	Age Group											Total
		0-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85+	
Severe	Male	53	111	122	230	218	150	86	61	31	5	1	1,068
Severe	Female	0	0	1	3	0	1	0	0	0	0	0	5
<b>Severe</b>	<b>Total</b>	<b>53</b>	<b>111</b>	<b>123</b>	<b>233</b>	<b>218</b>	<b>151</b>	<b>86</b>	<b>61</b>	<b>31</b>	<b>5</b>	<b>1</b>	<b>1,073</b>
Moderate	Male	6	14	29	50	47	47	40	37	17	15	1	303
Moderate	Female	0	0	0	1	0	0	0	0	0	0	0	1
<b>Moderate</b>	<b>Total</b>	<b>6</b>	<b>14</b>	<b>29</b>	<b>51</b>	<b>47</b>	<b>47</b>	<b>40</b>	<b>37</b>	<b>17</b>	<b>15</b>	<b>1</b>	<b>304</b>
Mild	Male	27	48	76	206	247	277	196	242	198	100	56	1,673
Mild	Female	0	1	5	13	20	21	22	15	7	2	1	107
Mild	Unknown	0	0	2	0	0	0	0	0	0	0	0	2
<b>Mild</b>	<b>Total</b>	<b>27</b>	<b>49</b>	<b>83</b>	<b>219</b>	<b>267</b>	<b>298</b>	<b>218</b>	<b>257</b>	<b>205</b>	<b>102</b>	<b>57</b>	<b>1,782</b>
Unknown	Male	0	1	0	3	3	1	2	3	1	0	2	16
Unknown	Female	0	0	1	4	8	4	5	4	2	1	0	29
<b>Unknown</b>	<b>Total</b>	<b>0</b>	<b>1</b>	<b>1</b>	<b>7</b>	<b>11</b>	<b>5</b>	<b>7</b>	<b>7</b>	<b>3</b>	<b>1</b>	<b>2</b>	<b>45</b>
Not Applicable	Female	0	0	0	2	1	2	2	2	0	0	0	9
<b>Not Applicable</b>	<b>Total</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>2</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>9</b>
<b>Total</b>	<b>-</b>	<b>86</b>	<b>175</b>	<b>236</b>	<b>512</b>	<b>544</b>	<b>503</b>	<b>353</b>	<b>364</b>	<b>256</b>	<b>123</b>	<b>61</b>	<b>3,213</b>

## Factor FVIII Deficiency - Patients with an active inhibitor, Statistics Canada age groups

Coverage: All of Canada Excluding British Columbia

Data source: CBDR

Year: 2020

All patients with an active inhibitor during 2020. Defined as a new inhibitor in 2020 or the most recent status from prior two years was positive at start of the year.

Severity	GenderCode	Age Group										Total
		0-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	
Severe	Male	10	4	10	14	6	3	7	4	4	1	63
<b>Severe</b>	<b>Total</b>	<b>10</b>	<b>4</b>	<b>10</b>	<b>14</b>	<b>6</b>	<b>3</b>	<b>7</b>	<b>4</b>	<b>4</b>	<b>1</b>	<b>63</b>
Moderate	Male	0	0	0	1	0	1	0	1	1	2	6
<b>Moderate</b>	<b>Total</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>6</b>
Mild	Male	0	0	1	1	1	0	2	3	0	0	8
<b>Mild</b>	<b>Total</b>	<b>0</b>	<b>0</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>0</b>	<b>2</b>	<b>3</b>	<b>0</b>	<b>0</b>	<b>8</b>
<b>Total</b>	<b>-</b>	<b>10</b>	<b>4</b>	<b>11</b>	<b>16</b>	<b>7</b>	<b>4</b>	<b>9</b>	<b>8</b>	<b>5</b>	<b>3</b>	<b>77</b>

## Factor FVIII Deficiency - Pediatric and Adult age groups

Coverage: All of Canada

Data source: CBDR, BC iChip

Year: 2020

Note: Inhibitor patients are included in this table

GenderCode	Severity	Age Group		Total
		under 18	18 and over	
Male	Severe	352	716	1,068
Male	Moderate	58	245	303
Male	Mild	214	1,459	1,673
Male	Unknown	1	15	16
<b>Male</b>	<b>Total</b>	<b>625</b>	<b>2,435</b>	<b>3,060</b>
Female	Severe	3	2	5
Female	Moderate	1	0	1
Female	Mild	8	99	107
Female	Unknown	2	27	29
Female	Not Applicable	0	9	9
<b>Female</b>	<b>Total</b>	<b>14</b>	<b>137</b>	<b>151</b>
Unknown	Mild	2	0	2
<b>Unknown</b>	<b>Total</b>	<b>2</b>	<b>0</b>	<b>2</b>
<b>Total</b>	<b>-</b>	<b>641</b>	<b>2,572</b>	<b>3,213</b>